

A taxonomic review of British decapod Crustacea

Joe S. H. Collins

8 Shaw's Cottages, Perry Rise, London, SE23 2QN, U. K. & Department of Palaeontology,
The Natural History Museum, Cromwell Road, London, SW7 5BD, U. K.

Abstract

Recent changes in the taxonomic position of a number of British decapod crustaceans are brought together from their respective publications. A new crab genus, *Stintonius*, is described to contain *Portunites subovata* Quayle & Collins, 1981, *Panopeus kemp* Ouayle & Collins, 1981, is also transferred to a new genus, *Sereneopeus*, and a subspecies, *Dromilites lamarckii humerosus* Quayle & Collins, 1981, is raised to specific status.

Key words: Crustacea, Decapoda, Mesozoic, Cenozoic, England

Introduction

Accounts of the fossil crabs of the British succession are widely distributed in the scientific literature and there is no recent review that considers their taxonomic diversity. Continuing research and refinement of diagnoses, often as not reflecting the constant differences of opinion between the 'splitters' and the 'lumpers', latterly largely attendant upon the revision of the *Treatise on Invertebrate Palaeontology* (Decapoda, Part R) (Feldmann, R. M. & Schweitzer, C. E. in progress), has led to the introduction of numerous changes in the taxonomic position of many species, a number of which effect the British fossil fauna dating back to the monographic work of Thomas Bell (1858).

Alterations and additions subsequent to such major, and other, publications, scattered among a diversity of journals and monographs, are herein gathered together, along with the description of a new genus, *Stintonius*, erected to accommodate a species left in limbo, while another species, left in similar state, is transferred to a new genus, *Sereneopeus*.

Lists of Plio/Pleistocene crabs published by Alfred Bell (1897, 1921) remain the sole contribution to our knowledge of the Quaternary species of the British Isles. Of the twenty six species listed in these two works all but two, *Cancer deshayesii* A. Milne Edwards, 1861, and *Maja verrucosa* H. Milne Edwards, 1834, are extant in British waters; sixteen have undergone taxonomic reform.

Coeloma sp., from Boxstones, in Suffolk requires verification, while A. Bell (1921) himself, cast doubt on the identification of *Calappa* sp. that he first recorded in 1897. *Cancer deshayesii* is considered herein; *Maja verrucosa*, recorded from an almost entire carapace from the Coralline Crag of Butley, presently occurs in the Mediterranean (Zariquiey Alvarez, 1946). *The Monograph of the Crustacea of the London Clay* by Thomas Bell (1858) remains the only collective work concerning the Eocene species. Additions, largely concerning new species from the neighbourhood of Portsmouth, were made by Woodward (1867, 1871, 1873); *Plagiolophus wetherelli* Bell, 1858 was properly referred to as [*Clyphithyreus*=] *Glyphithyreus wetherelli* (Bell) by Brown & Castell (1954). Glaessner & Withers (1931) and Collins (1961) discussed Lower Eocene crabs. Cooper (1974) published a stratigraphical distribution of the English Palaeocene decapods, in which he included *Harpactoxanthopsis* cf. *quadrilobata* (Desmarest, 1822) from the Isle of Sheppey, based on an oral report by the present author. Some generic revision, largely attendant upon synonymy, was listed by Morris (1980) in which work the synonymy surrounding *Zanthopsis leachii* (Desmarest, 1822) and allied species was resolved. In the same work, *Xanthopsis leachii* var. Bell, 1858 (pl. 1, fig. 10) was properly identified as *Zanthopsis dufouri* (H. Milne Edwards, 1850). Crane & Quayle (1986) and Crane (1981) described new Middle Eocene species and discussed London Clay species. Quayle (1984) described a new species,

Portunites stintoni, from the London Clay, and in 1987 revised and added new macruran species from the London Clay, Bracklesham and Barton Beds. A contribution to Middle and Upper Eocene crab species by Quayle & Collins appeared in 1981.

The description of a rare Upper Albian glypheiid, *Glypheopsis sanctaerucis woodsi* Collins, appeared in 1969, together with the description of the nominative species which comes from the Upper Albian of Annapol on the Vistula, Poland. Only three works concerning Cretaceous species have appeared since the major review of British Cretaceous crabs by Wright & Collins was published in 1972; Simpson (1983) recorded *Mithracites vectensis* Gould, 1859, *Callianassa* claws and, following Förster's (1971, see below) generic opinion, *Mecochirus magnus* (McCoy, 1849) from the Lower Aptian, Atherfield Clay, Punfield Marine Band of Dorset; but in 1985 he assigned *Mecochirus magnus* to a new genus, *Meyerella*, in a list of species from the Lower Aptian Atherfield Clay Formation. At the same time (Simpson, 1985) notice was drawn to the occurrence of *M. vectensis* from the Atherfield Clay of Sussex and Surrey, from which latter county *Hoploparia longimana* (G. B. Sowerby) was also recorded. Wright (1997) described a new galatheid and prosoponid, *Rathbunopon? atherfieldensis*, from the Upper Aptian of the Isle of Wight.

No collective work on British Jurassic decapods exists, although some attempt was made by Salter & Woodward (1865) (Woodward & Salter of some authors) to illustrate the known species at the time, and a comprehensive assessment of taxa was made by Van Straelen (1925). Full descriptions of all known macruran species were included in the monographic work of Woods (1925-1931). The earliest description in a British journal appears to be that of *Astacus leachii* Mantell, 1822, later (1849) assigned to *Enoploclytia* by McCoy. *Gebia clypeatus* Carter, 1898, was included in *Mecochirus* by Förster (1971). *Eryma portlandica* Woods (1930) was relegated to junior synonymy of *Eryma duretrei* Sauvage, 1891, by Förster (1966), who later (1971) included *Gebia clypeatus* Carter (1898) in *Mecochirus*. Species assigned to the Erymidae, Glypheidae, Mecochiridae and Palinuridae underwent considerable revision by Förster (notably in 1966, 1971, 1973).

In the discussion of their new species, *Pseudoglyphea foersteri*, from the Lower Jurassic (Pliensbachian) of Scotland, Feldmann *et al.* (2002) concluded that cephalic characters of the Mecochiridae were more consistent with

the Astacidae than the Palinura.

Diagnostic descriptions of the thalassinoids, *Etallonia isochela* (Woodward, 1876) (Upper Kimmeridgian) and *Magila pichleri* Opper, 1962 (Callovian) were given by Förster (1977), who at the same time questioned the status of *Magila laevimana* Carter, 1886, and *Magila dissimilis* Carter, 1886 (both Oxfordian). The description of the Oxfordian pagurid *Goniochirus cristatus* by Carter (1886) was followed (1898) by his recording of the galatheid *Gastrosachus wetzleri* von Meyer, 1854, from the Coral Rag.

The first notice of a British Jurassic crab, *Protocarcinus longipes* Bell MS, was that figured by Salter & Woodward (1865) and described as *Palaeinachus longipes* by Woodward (1866). However, Woodward was obviously unaware that he had published a junior synonym of *Homolus audini* Deslongchamps, 1835, a fact made known by Glaessner (1929a), by which time *audini* had been assigned to *Protocarcinus* by Van Straelen (1923). A further change of name was effected by Wehner (1988) when she transferred the species to *Foersteria* Wehner, 1988. The description of *Prosopon mammillatum* by Woodward (1868) was followed by that of *Prosopon richardsoni* Woodward, 1907. Descriptions of both species were extended by Withers (1951), who, previously (1932) had described the earliest known British crab, *Eocarcinus praecursor* from the Upper Lias of Oxfordshire – a species which received further attention from Förster (1979). Donovan (1962) described *Pithonoton* sp. and extended the geographic range of *P. richardsoni*, and the prosoponids received further attention from Wehner (1988). Contributions were made to British Jurassic decapods by Glaessner (1933) and Fraaye & Collins (1996) described two species from the Portland beds of Dorset. However, from the generic point of view, the galatheid, *Eomunidopsis portlandica*, included by Fraaye and Collins (1996) in the Munidopsinae Ortmann, 1898, was transferred to the Galatheinae Samouille, 1819, by Schweitzer & Feldmann (2000c).

Tropifer laevis Gould, 1857, from the Upper Triassic Rhaetian stage of Avon (formerly Gloucestershire) was redescribed and tentatively placed in the Coleiidae by Duffin (1978).

Systematics

Infraorder Anomura MacLeay, 1838
Superfamily Thalassinoidea Latreille, 1831

Family Callianassidae Dana, 1852

Genus *Callianassa* Leach, 1814 *sensu lato*

"*Callianassa*" *batei* Woodward, 1869

1869 *Callianassa Batei* Woodward, 75, pl. 2, fig. 4.

Remarks: This species, founded on chelae from the Headon and Hampstead Beds, Upper Eocene/Lower Oligocene, of the Isle of Wight, and recorded only by figured left and right chelae, is a senior homonym of the Recent *Callianassa batei* Borradaile (1903, p. 546), a species with an involved history which included the synonym and replacement name *Callianassa profunda* Biffar, 1973 (Sakai, 1999).

Family Galatheididae Samouelle, 1819

Subfamily Galatheinae Samouelle, 1819

Genus *Luisogalatea* Karasawa & Hayakawa, 2000

?*Luisogalatea* sp. Wright, 1996

1996 *Galathea* sp., Wright, 137, fig. 8.

Remarks: Reexamination of the specimen, from the Atherfield Clay, Lower Greensand, Crackers Bed, Atherfield Point, Isle of Wight, showed it to be an internal cast with remains of shell thickness adpressed to the surrounding matrix, thus, evidence of possible lateral spines – a generic requirement – is largely illusory. Otherwise, the specimen agrees with *Luisogalatea tomatai* Karasawa & Hayakawa, 2000, in having a long triangular, smooth margined rostrum without a median ridge; the gastric region is rather more granular overall, rather than limited to the hepatic regions as in *L. tomatai*; the cervical and postcervical furrows are equally well developed and the branchial region is similarly transversely ridged.

The presence of a smooth sided, non-serrate rostrum and lack of gastric ridges immediately distinguishes *Luisogalatea* from *Galathea*, and it would seem that the present carapace more readily suits the requirements for *Luisogalatea*. Thus, it considerably extends the known geological and geographical ranges of the genus, hitherto known from the Upper Cretaceous Yezo Group of Japan and possibly by *Luisogalatea cobbani* (Bishop, 1985) from the Campanian of Colorado, which was included in the genus by Karasawa & Hayakawa (2000), but not referred to by Schweitzer & Feldmann (2002c).

Superfamily Paguroidea Latreille, 1802

Family Paguridae Latreille, 1802

Genus *Pagurus* Fabricius, 1775

***Pagurus bernhardus* Linnaeus, 1758**

Remarks: This species was recorded by A. Bell (1921) from chelae fragments from the Coralline Crag, Red Crag and Pleistocene from several localities, as *Eupagurus*, a genus recognised as a junior synonym of *Pagurus* (q.v. Glaessner, 1969).

Section Dromiacea de Haan, 1833

Superfamily Dromioidea de Haan, 1833

Family Prosopidae von Meyer, 1860

Genus *Rathbunopon* Stenzel, 1945

***Rathbunopon?* *atherfieldensis* Wright, 1997**

Remarks: This species may well be conspecific with the *Rathbunopon* referred to by Simpson (1985). This possible new *Rathbunopon* is from the Lower Aptian of the Isle of Wight, and was considered by Wright (1997) to have, "some resemblance [of *R.? atherfieldensis*] to the fragmentary holotype of the Hauterivian" considerably extends the lower range of the genus from the Lower Albian. In the same work (1997), Wright drew attention to *Homolopsis tuberculata* Van Straelen, 1936, from the Hauterivian which may also be a *Rathbunopon*.

Family Dromiidae de Haan, 1833

Genus *Basinotopus* McCoy, 1849

Type species: *Inachus lamarckii* Desmarest, 1822, by monotypy.

***Basinotopus lamarckii* (Desmarest, 1822)**

1822 *Inachus Lamarckii* Desmarest, 116.

1849 *Basinotopus lamarckii* (Desmarest); McCoy, 168.

1858 *Dromilites Lamarcki* (Desmarest); Bell, 29.

1981 *Dromilites lamarckii lamarckii* (Desmarest); Quayle & Collins, 738.

In prep. *Basinotopus lamarckii* (Desmarest); Collins & Jakobsen.

Remarks: While superficially close, *Basinotopus lamarckii* is distinguished from *Dromilites bucklandii* H. Milne Edwards, 1837, the type species of *Dromilites*, in having a produced rostrum between elongated spines (rostral horns) extending from the upper orbital margin. These characters are invariably abraded and reconstructed figures/retouched photos of *D. lamarckii* have hitherto shown only a short, triangular rostrum. However, the rostral characters, vaguely referred to by Bell (1858) and remarked upon by Glaessner (1929b)

when discussing *Dromilites alpina* Glaessner, 1929b, are revealed in a recently developed carapace (In48212) from Bognor Regis, in The Natural History Museum London (BMNH), and a fragmentary carapace from the Isle of Sheppey, in the Geological Museum, University of Copenhagen, which retains the rostrum, thereby allying *B. lamarckii* with an as yet, unpublished species from the Middle Eocene of Denmark.

Genus *Dromilites* H. Milne Edwards, 1837

Type species: Dromilites bucklandii H. Milne Edwards, 1837, by monotypy.

***Dromilites humerosus* Quayle & Collins, 1981**

1981 *Dromilites lamarckii humerosus* Quayle & Collins, 731.

Remarks: This species, founded on three fragmentary carapaces from the Elmore Formation, Bracklesham Group, of Lee-on-the-Solent, Hampshire, is readily distinguished from *Basinotopus lamarckii* by its prominent subhepatic lobes. Furthermore, although the front is poorly preserved on all three available specimens, the holotype has a short, weakly sulcate rostrum, bounded by continued thickening of the upper orbital margin. No basal scars can be discerned to support presence of rostral horns. Therefore, Quayle and Collins's (1981) assumption of derivation of *D. l. humerosus* from *D. l. lamarckii* is illusionary, and the former species is here raised to specific status and retained, for the time being, in *Dromilites*.

Superfamily Homoloidea de Haan, 1833

Genus *Mithracites* Gould, 1859

***Mithracites vectensis* Gould, 1859**

Remarks: Described within the Cyonomidae by Wright & Collins (1972), this species was considered by Guinot & Tavares (2001) to have podotreme sexual characters more closely related to the Homoloidea and transferred the genus to that superfamily without reference to a family.

Section Podotremata Guinot, 1977

Family Etyidae Guinot & Tavares, 2001

Genera *Etyus* Leach (in Mantell, 1822) :

***Xanthosia* Bell, 1863**

Remarks: Discovery of a carapace (B22703) of *Etyus martini* Mantell, 1822, retaining 'openings of oviducts' on misinterpreted 3rd sternites, played an influential part in

restoring *Etyus* and the closely allied genus *Xanthosia*, from the Dynomenidae (q.v. Glaessner, 1969) to the Xanthidae by Wright & Collins (1972). Recent research by Guinot & Tavares (2001) on the presence of spermathecae in diverse families and genera, embraced *Etyus* and *Xanthosia*. The above mentioned pores were recognised as spermathecae opening between the 7th/8th sternal sutures. This a character which, while not conforming to the Xanthidae, is shared within the Podotremata, thus warranting placement in a new family.

Section Eubrachyura Saint Laurent, 1980

Subsection Raninoidea de Haan, 1839

Superfamily Raninoidea de Haan, 1841

Family Raninidae de Haan, 1841

Subfamily Palaeocorystinae Lőrenthey in Lőrenthey & Beurlen, 1929

Type genus: Palaeocorystes Bell, 1863 (= *Notopocorystes* McCoy, 1849).

Remarks: Palaeocorystinae is the valid name for this subfamily under Article 40.1 of the International Code of Zoological Nomenclature (1999, 4th ed., p. 46). The action of Haj & Feldmann (2002) is incorrect either under Article 40.1, or 40.2 of the code had their work been published before 1961.

Genera *Notopocorystes* McCoy, 1849:

***Cretacorantina* Mertin, 1941:**

***Eucorystes* Bell, 1863**

1972 Wright & Collins, 73-86, pl. 13, figs. 4-6; pl. 14, figs. 1-7; pl. 15, figs. 1-4; pl. 17, figs. 1-5; pl. 18, figs. 1-5; pl. 2, fig. 7; pl. 22, fig. 7. (See also for prior synonymy.)

Remarks: Treated as subgenera by Wright & Collins (1972), with *Notopocorystes* as nominate genus, the three abovementioned Albian – Lower Chalk genera were raised to full generic status by Tucker (1998), who at the same time, raised the three subspecies, of the nominate genus, *N. stokesii stokesii* Mantell, 1844, *N. stokesii praecox* Wright & Collins, 1972 and *N. stokesii serotinus* Wright & Collins, 1972, to specific rank. However, since the phylogenetic transition of these forms is readily traceable, absolute distinction becomes finite only with advanced development. With specific status, progressive forms would be left in limbo, therefore, retention of subspecific status would seem the more appropriate. The subspecies of *Eucorystes carteri* (McCoy, 1854), *E. c. ligulatus* Wright & Collins, was overlooked by Tucker (1998).

Functional morphology of the cuticular structure present in *Eucorystes [carteri] carteri*, (McCoy, 1854) was discussed and figured by Haj & Feldmann (2002).

Superfamily Cyclodorippoidea Ortmann, 1892

(= Superfamily Tymolidae Alcock, 1896)

Family Torynommatidae Glaessner, 1980

(*rectae*, ICZN, from Torynommatidae Glaessner, 1980)

Genus *Dioratiopus* J. Woods, 1953

Remarks: Some British members of this genus have had an extremely chequered taxonomic history. *Dioratiopus spinosus* (Van Straelen, 1936) and *Dioratiopus depressus* (Carter, 1898), both from the Albian, were considered by their respective authors to belong to *Homolopsis*. In the absence of diagnostic *lineae homolicae*, those taxa, with other, newly described, species (Lower Aptian-Cenomanian) were placed in *Glaessneria* by Wright & Collins (1972), but that genus, found to be a homonym of *Glaessneria* Takeda & Miyake, 1969, was replaced by *Glaessnerella* Wright & Collins, 1975 (family Cymonomidae Bouvier, 1898). Transferred to *Dioratiopus* and considered close to *Torynomma* J. Woods, 1953, Glaessner (1980) included both genera in a new family. The earliest known (Jurassic) member, *Dioratiopus primitivus* from the Portland Beds of Dorset, was described by Fraaye & Collins (1996), who remarked upon the relationship of *Dioratiopus* to the prosoponid, *Foersteria*.

Superfamily Calappoidea de Haan, 1833

Family Necrocarcinidae Förster, 1968b

Genus *Campylostoma* Bell, 1858

Remarks: Previously contained in the Calappidae, the genus was transferred to the Necrocarcinidae by Schweitzer & Feldmann (2000b). The full extent of the spine at the lateral angle – referred to by Bell (1858) as 'evidently unnaturally developed' – of the sole species *Campylostoma matutiforme* Bell, 1858 (London Clay), was figured by Collins (1961). As figured in Schweitzer & Feldmann (2000b), the reconstruction is similar to that presented by Salter & Woodward (1865).

Genus *Goniochele* Bell, 1858

***Goniochele angulata* Bell, 1858**

Remarks: Previously classified in the Dorippoidea, Dorippinae, this London Clay species was transferred to the Necrocarcinidae by Schweitzer & Feldmann (2000b).

Genus *Orithopsis* Carter, 1872

***Orithopsis tricarinatus* (Bell, 1863) nov. comb.**

1972 *Necrocarcinus tricarinatus* Bell, 1863; Wright & Collins, 1972, 66. (See also for previous synonymy).

Remarks: Recognised as being conspecific, *Orithopsis bonneyi* Carter, 1872, was classified by Wright & Collins (1972) as a junior synonym of *Necrocarcinus tricarinatus* Bell, 1863. On raising the Necrocarcininae Förster, 1968, to full family status, Schweitzer & Feldmann (2000a) believed that *Orithopsis* should be kept distinct. This, being accepted, *tricarinatus* Bell, 1863, remains the senior taxon in the new combination *Orithopsis tricarinatus* (Bell, 1863) and in this form should replace '*Orithopsis bonneyi* Carter, 1872' elected as type species by Schweitzer & Feldmann (2000b).

Subgenera *Paranecrocarcinus*

(*Paranecrocarcinus*) Van Straelen, 1936:

Paranecrocarcinus (Pseudonecrocarcinus)

Förster, 1968b

Remarks: These subgenera were distinguished by Wright & Collins (1972), largely on the presence or [presumed] absence of post-rostral slits; forms in *Pseudonecrocarcinus* having 1 or 2 pairs of slits, while none was present in [available] species assigned to *Paranecrocarcinus*. However, as shown by Wright (1997), one pair of slits is present in the type species, *Paranecrocarcinus hexagonalis* (Van Straelen, 1936) (Cenomanian, France), and one pair of slits is seen in well preserved specimens of *Paranecrocarcinus (Paranecrocarcinus) digitatus* Wright & Collins, 1972. Maintenance of subgenera was questioned by Wright, who was in favour of abandonment of the junior *Pseudonecrocarcinus*.

Superfamily Cancroidea Latreille, 1802

Family Atelycyclidae Ortmann, 1893

Genus *Atelycyclus* Leach, 1814

***Atelycyclus rotundatus* (Olivi, 1792)**

Remarks: Recorded from the Coralline Crag of Aldeburgh, Suffolk, by A. Bell (1897) as *Atelycyclus heterodon* Leach, 1815, this extant species was recognised as a junior synonym of *A. rotundatus* by Ingle (1980). *Cancer (Hippra) septemdentatus* Montague, 1813 is also placed in junior synonymy by Ingle (1980), as it was by Adema (1991), who included the earlier date of *Ateleycyclus septemdentatus* Montague, 1808, and Salva & Feldmann

(2001).

Family Cancridae Latreille, 1802
 Subfamily Lobocarcininae Beurlen, 1930
Genus *Lobocarcinus* Reuss, 1857

***Lobocarcinus sismondai* (von Meyer, 1859)**

1861 *Cancer deshayesii* A. Milne Edwards, 74.

Remarks: Formerly described and figured as *Cancer deshayesii* by A. Bell (1921), a partial carapace from the Pliocene Coralline Crag of Aldeburgh, Suffolk now in the York Museum (No. YM68), retains entire left 6/7th typically spinulose marginal lobes conforming to figures in Bonfiglio & Donadeo (1982). This species, also recorded in the fossil record by Holthuis (1949) on the evidence of isolated dactyli and fixed fingers from Miocene-Lower Pleistocene deposits in The Netherlands, was placed by Glaessner (1924) in *Lobocarcinus* Reuss, 1857, as a subgenus of *Cancer*, but was soon returned to *Cancer*. *Cancer deshayesii* was synonymized with *Cancer sismondai* by Glaessner (1929a). Müller (1984) considered that there is no difference between *Cancer illyricus* Bittner, 1883, and *C. sismondai* which was transferred to *Lobocarcinus* Reuss, 1857, by Schweitzer & Feldmann (2000a). Neither the British nor Dutch records for this species were mentioned specifically by Schweitzer & Feldmann (2000a), who simply recorded northern Europe, with an upward range limited to the Pliocene.

Family Cheiragonidae? Ortmann, 1893

***Stintonius* gen. nov.**

Type species: *Portunites subovata* Quayle & Collins, 1981, by monotypy.

Diagnosis: Carapace subovate with four slender anterolateral spines increasing in length posteriorly, regions well defined and weakly tumid; epibranchial lobe weakly ridged; a tubercle on each mesobranchial and two on each metabranchial lobe form a diverging row, and one marginal tubercle on each metabranchial lobe.

Derivation of name: In memory of the late F. C. Stinton, an outstanding authority on the Eocene Barton Beds of Hampshire.

***Stintonius subovatus* (Quayle & Collins, 1981)
 nov. comb.**

1981 *Portunites subovata* Quayle & Collins, 749.

Remarks: In a revision of *Portunites* by Schweitzer & Feldmann in 1999 (*vide* Schweitzer & Feldmann, 2000d),

subovata was found to have characters inconsistent with that genus and removed therefrom. No further steps were taken, apart from suggesting the species might be referred to *Montezumella* Rathbun, 1930b, a genus represented in the Barton fauna by *Montezumella scabra* Quayle & Collins, 1981. Indeed, there is remarkable conformation of the carapace outline of *subovatus* with that of *Montezumella*. However, with the disparity in size between the two species, it is difficult to conceive the degree of ontogenetic development necessary for *subovata* to develop the characters of *M. scabra*, and inclusion in a new genus, tentatively included in the Cheiragonidae, is considered necessary. For description and figures, see Quayle & Collins (1981).

Superfamily Portunoidea Rafinesque, 1815

Family Portunidae Rafinesque, 1815

Subfamily Polybiinae Ortmann, 1893

Genus *Leiocarcinus* Stimpson, 1870

Remarks: The following species recorded from Pliocene, Pleistocene or Holocene deposits were all listed in *Portunus* by A. Bell, 1897: *P. corrugatus* Pennant, 1777; *P. depurator* Linnaeus, 1758; *P. puber* Linnaeus, 1767 and *P. pusillus* Leach, 1815. In 1921, A. Bell listed a further two species, *P. marmoreus*, Leach, 1814 and *P. ?holsatus* (Fabricius, 1798). All were included in *Leiocarcinus* by Ingle (1980) in his survey of Recent species. Adema (1991) recognised, *L. holsatus*, *L. marmoreus* and *L. pusillus* as present in the Recent Dutch and Belgian fauna, but recorded *puber* in *Necora* Holthuis, 1987.

Superfamily Xanthoidea MacLeay, 1838

Family Goneplacidae MacLeay, 1838

**Genus *Orthakrolophus* Schweitzer & Feldmann,
 2001**

***Palaeograpsus bartonensis* Quayle & Collins, 1981:
Palaeograpsus depressus Quayle & Collins,
 1981**

Remarks: Both of these species, considered to have characters inconsistent with *Palaeograpsus*, were assigned, to *Orthakrolophus* by Schweitzer & Feldmann, 2001, with *depressus*, from the Bracklesham Group of Lee-on-the-Solent and Barton Beds of Christchurch Bay and the Isle of Wight, as the type species. *Orthakrolophus bartonensis* (Quayle & Collins) is known only from Barton Beds of Christchurch Bay.

Family Panopeidae Ortmann, 1893

Genus *Sereneopeus* gen. nov.

Type species: Panopeus kemp Quayle & Collins, 1981, by monotypy.

Diagnosis: Carapace hexagonal with rounded anterolateral margins; wider than long; front gently sinuous with shallow median notch; first, of four, anterolateral spines weak, third and fourth spines oblong rather than triangular, lobes weakly defined; epibranchial lobe weakly ridged.

Derivation of name: In recognition of the contributions to the study of crabs made by Raoul Serène + familial root.

Sereneopeus kemp

(Quayle & Collins, 1981) nov. comb.

1981 *Panopeus kemp* Quayle & Collins, 752.

Remarks: Considered to have characters inconsistent with *Panopeus*, *kemp* was removed from that genus by Schweitzer (2000). However, in general diagnostic characters, *kemp* complies with the requirements of Panopeidae, yet which, in combination, are distinct from genera presently included in that family. The principle reason for removal of *kemp* from *Panopeus* was lack of, "transverse [hepatic, epigastric, protogastric] ridges, fewer, more poorly developed spines and carapace regions of a different shape'. It should be borne in mind that all available specimens of *kemp* are internal casts; 'shell'-surface detail is absent – and surface detail is frequently not reflected in internal casts. Nevertheless, *kemp* has the, "five marginal spines, the first forming the outer angle of the orbit" (Quayle & Collins, 1981), a carapace that is wider than long, and weakly sinuous margins either side of a frontal notch, thus far complying with the familial requirement. While lacking the abovementioned transverse ridges (and in this respect resembling the somewhat aberrant *Panopeus whittenensis* Glaessner, 1980, Upper Eocene, Australia), the epibranchial lobe is as strongly ridged as in the Recent, *e.g.* *Panopeus purpureus* Lockington, 1877, which also has comparatively weakly defined lobes. These conditions are particularly noticeable among species of *Eurypanopeus* A. Milne Edwards, 1880, in which genus the development of the anterolateral spines is variable (*vide* Rathbun, 1930a).

Sereneopeus also shares characters seen in the Piummidae Samouelle, 1819, but can be eliminated from that family by its greater length/width ratio, in having a relatively straight, unadorned frontal margin and large,

triangular hepatic regions.

Family Carcineretidae Buerlen, 1930

Genus *Withersella* Wright & Collins, 1972

***Withersella crepitans* Wright & Collins, 1972**

Remarks: Vega *et al.* (1997) were adamant in their exclusion of *Withersella crepitans* in the Carcineretidae Buerlen, 1930, in which family it was placed by Wright & Collins (1972). Glaessner's (1980) reference of *Withersella* to the Tymolidae was reiterated. However, Wright (1997) provided additional details of the front of *W. crepitans* showing that it is, "extremely close to that of *Carcineretes walcotti* Withers [Maastrichtian, Jamaica] except for the greater projection of the rostrum of *Withersella*." Taking into account the considerable difference in age between the species retained in Carcineretidae by Vega *et al.* (1997), their arguments against inclusion of *Withersella* within the Carcineretidae – that "*Withersella* lacks a transverse ridge, has more distinctly defined surface regions and is square rather than wider than long" - provides no latitude for possible phylogenetic development within the family. A more important character, that is, the possession of a flattened, paddle-shaped fifth limb, has yet to be determined in *Withersella* and ultimate familial position is probably best deferred until this character is established.

The transverse ridge (across the protogastric lobes) of *Carcineretes planetarius* Vega *et al.* (1997) (Lower Maastrichtian of Belize) is weaker than that of the type species, *Carcineretes woolcotti*, which, while lacking the definition of the anterior mesogastric process seen in *C. planetarius*, has regions as well defined as *Withersella*.

Unfortunately, Wright did not draw attention to a massive bopyric swelling occupying the entire right metabranchial region of his figured specimen (1997, fig. 12) which comes from the Lower Greensand, Crackers Bed, Lower Aptian of the Isle of Wight.

Acknowledgements

Warmest thanks are extended to Dr Phil Manning (Yorkshire Museum) for loan of *Lobocarcinus sismondai*, to Dr S. K. Donovan (Nationaal Natuurhistorisch Museum, Leiden), and Dr Hiroaki Karasawa (Mizunami Fossil Museum), for their encouragement and critical comments on the MS, and to Mr John Cooper (The Natural History Museum, London) for assistance with

references. Verification of the status of Palaeocorystinae was kindly made by Dr A. Wakeham-Dawson, Executive Secretary, International Code of Zoological Nomenclature (The Natural History Museum, London).

References

- Adema, J. P. H. M. (1991), *De Krabben van Nederland en België*. Nat. Natuurh. Mus. Leiden, xii+244 p.
- Alcock, A. (1896), The Brachyura Oxystoma: Materials for a Carcinological Fauna of India. No. 2. *Jour. Asiat. Soc. Bengal*, **68** (II:3), 123-169.
- Bell, A. (1897), A synopsis of the Crustaceans and Echinodermata of the Upper Tertiaries. *Rpt. Yorks. Philos. Soc. for 1896*, 12 p.
- Bell, A. (1921), Notes on the later Tertiary. *Rpt. Yorks. Philos. Soc. for 1920*, 21 p.
- Bell, T. (1853), *A History of the British Stalk-eyed Crustacea*. John van Voorst, London. Ixii+386 p.
- Bell, T. (1858), *A monograph of the fossil malacostracous Crustacea of Great Britain. Part I. Crustacea of the London Clay*. Palaeontogr. Soc. [Monogr], London, 44 p.
- Bell, T. (1863), *Idem*. Part II. Crustacea of the Gault and Greensand. *Ibid.*, viii+40 p.
- Beurlen, K. (1930), Vergleichende Stammesgeschichte, Grundlagen Methoden, Probleme unter besonderer Berücksichtigung der höheren Krebse. *Fortschr. Geol. Palaeont.*, **8**, 317-586.
- Bishop, G. A. (1985), A new crab, *Eomunidopsis cobbani* n. sp. (Crustacea, Decapoda), from the Pierre Shale (Early Maastrichtian) of Colorado. *J. Paleont.*, **59** (3), 601-604.
- Biffar, T. A. (1973), The taxonomic status of *Callianassa occidentalis* Bate, 1888, and *C. Batei* Borradaile, 1903 (Decapoda, Callianassidae). Biological results of the University of Miami Deep-Sea Expeditions **98**, *Crustaceana*, **24** (2), 224-230.
- Bittner, A. (1883), Neue Beiträge zur Kenntniss der Brachyuran Fauna des Alttertiärs von Vinzenta und Verona. *Denskr. K. Akad. Wissensch., Wien*, **46**, 229-316.
- Bonfiglio, L. and G. Donadeo (1982), *Cancer sismondai* Meyer nel Pliocene di Torre dell'Orso (Puglia). *Atti Soc. Ital. nat. Museo. civ. Stor. nat. Milano*. **123** (2-3), 255-296.
- Borradaile, L. A. (1903), Marine Crustaceans. Pts. IV – VII. IV. Some remarks on the classification of the crabs. In Gardiner, J. S., *The Fauna and Geography of the Maldive and Lacadive Archipelagoes ...* **14**, 424-442. Cambridge.
- Brown, M. M. and C. P. Castell (1954), The London Clay of Oxshott, Surrey. *London Naturalist* (1953), no. 86, 61-64.
- Bouvier, E. -L. (1898), Sur la Xanthes des mers d'Europe. *Feuil. Jeun. Nat.*, 3^e sér., **28**, no. 332, 133-137.
- Carter, J. (1872), On *Orithopsis Bonneyi*, a new fossil crustacean. *Geol. Mag.*, **9**, 529-532.
- Carter, J. (1886), On the decapod Crustaceans of the Oxford Clay. *Q. Jl. Geol. Soc. Lond.*, **42**, 542-559.
- Carter, J. (1898), A contribution to the palaeontology of the decapod Crustacea of England. *Q. Jl. Geol. Soc. Lond.*, **54**, 15-44.
- Collins, J. S. H. (1961), Eocene crabs in a London Clay nodule. *Palaeontology*, **4** (1), 85-86.
- Collins, J. S. H. (1969), Some decapod crustaceans from the Lower Cretaceous of Poland and England. *Acta palaeont. Polon.*, **19** (4), 564-574.
- Collins J. S. H. and S. L. Jakobsen (in progress), Crabs (Crustacea, Decapoda) from the Middle Eocene (Lutetian) Lillebælt Clay Formation of southeast Jutland.
- Cooper, J. (1974), The stratigraphical distribution of the English Palaeogene Decapoda. *Tertiary Times*, **2** (2), 83-85.
- Crane, M. D. (1981), Hexapod crabs of the genus *Goniocypoda* H. Woodward from the Upper Eocene of Hampshire. *Zool. Journ. Linn. Soc.*, **72**, 1-19.
- Crane, M. D. and W. J. Quayle (1986), Two new hexapod crabs of the genus *Goniocypoda* Woodward (Crustacea, Decapoda) from the Hampshire Basin. *Tert. Res.*, **7** (4), 101-110.
- Dana, J. D. (1852), Crustacea. In *U.S. Exploring Exped. during the years 1838, 1839, 1840, 1841, 1842 under the command of Chas. Wilkes, U. S. N.*, 13 (Crustacea, pt. 1), viii+685 p.
- Deslongchamps, E. Eudes- (1835), Mémoire pour servir à l'histoire naturelle des Crustacés fossiles *Mém. Soc. Linn. Normandie*, **5**, 37-46.
- Desmarest, A.-G. (1822), *Histoire naturelle des Crustacés fossiles. Les Crustacés proprement dits*. Paris, 116 p.
- Donovan, D. T. (1962), New examples of the fossil crab *Pithonoton* from the Inferior Oolite of Gloucestershire. *Proc. Geol. Assoc.*, **73** (2), 193-196.
- Duffin, R. C. (1978), *Tropifer laevis* Gould (Coleiidae: Crustacea) from the Rhaetian Bone Bed of Aust Cliff, Avon. *Zool. J. Linn. Soc.*, **64**, 177-185.
- Fabricius, J. C. (1775), *Systema Entomologiae sistens Insectorum classes, ordines, genera, species, adjectis synonymis, locis, descriptionibus*. Flensburgi et Lipsiae, 832 p.
- Fabricius, J. C. (1798), *Entomologia systematica emendata et aucta ... adjectis synonymis, locis, observationibus, discriptionibus. Supplement. Hafniae*, 572 p.
- Feldmann, R. M., G. Crisp, and D. Pirrie (2002), A new species of glypheoid lobster, *Pseudoglypheo foersteri* (Decapoda: Astacidea: Mecochiridae) from the Lower Jurassic (Pliensbachian) of Raasay, Inner Hebrides. *Palaeontology*, **45** (1), 23-32.
- Feldmann, R. M., A. C. Ocampo, and K. O. Pope (1997), A new species of Late Cretaceous crab (Brachyura: Carcineretidae) from Albion Island, Belize. *J. Paleont.*, **71** (4), 615-620.
- Feldmann, R. M. and C. E. Schweitzer (in progress). *Treatise on Invertebrate Paleontology*, Pt. R.
- Förster, R. (1968), Über die Erymiden, eine alte konservative

- Family der mesozoischen Dekapoden. *Palaeontogr.*, **125A**, 61-175.
- Förster, R. (1971), Die Mecochiridae, eine spezialisierte, Familie der mesozoischen Glyphoidea (Crustacea, Decapoda). *N. Jb. Geol. Paläont. Abh.*, **137**, 396-421.
- Förster, R. (1973), Untersuchungen an oberjurassischen Palinuridae (Crustacea, Decapoda). *Mitt. Bayer. Staatssaml. Paläont. Hist. Geol.*, **13**, 31-46.
- Förster, R. (1977), Untersuchungen an jurassischen Thalassinioidea (Crustacea, Decapoda). *Mitt. Bayer. Staatsslg. Paläont. Hist. Geol.*, **17**, 137-156.
- Förster, R. (1979), *Eocarcinus praecursor* Withers (Decapoda, Brachyura) from the Lower Pliensbachian of Yorkshire and the early crabs. *N. Jb. Geol. Paläont., Mh.*, **1**, 15-27.
- Fraaye, R. F. and J. S. H. Collins (1996), Two new decapod crustaceans from the Portlandian of Dorset, England. *Proc. Geol. Assoc.*, **107**, 323-326.
- Glaessner, M. F. (1924), Ueber eine neue miozäne Krabbe und die Brachyurenfauna des Wiener Beckens. *Veerhandl. Geol. Bundesanst. Wien*, **6**, 109-118.
- Glaessner, M. F. (1929a), *Fossilium Catalogus 1: Animalia, pars. 41 Crustacea decapoda*. Berlin, 464 p.
- Glaessner, M. F. (1929b), Decapodenfauna. *Neun. Jb. Miner. Geol. Paläont.*, Biel, Band 63, B, 137-176.
- Glaessner, M. F. (1933), Die Krabben der Jura-Formation. *Zbl. Mineral. Geol. Paläont.*, Abt. B, 178-199.
- Glaessner, M. F. (1969), Decapoda. In Moore, R. C. and Teichert, C. (eds.), *Treatise on Invertebrate Paleontology, Part R, Arthropoda. 4, 2*, R399-R533, R626-628. Geological Society of America and University of Kansas Press. Boulder & Lawrence.
- Glaessner, M. F. (1980), New Cretaceous and Tertiary crabs (Crustacea: Brachyura) from Australia and New Zealand. *Trans. R. Soc. S. Aust.*, **6**, 171-192.
- Glaessner, M. F. and T. H. Withers (1931), London Clay Crabs of the Family Raninidae. *Ann. Mag. nat. Hist.*, ser. 10, **8**, 484-493.
- Gould, C. (1857), On a new crustacean (*Tropifer laevis*. Gould) from the Lias bone bed. *Q. J. geol. Soc. Lond.*, **13**, 360-363.
- Gould, C. (1859), Description of a new fossil Crustacean from the Lower Greensand. *Q. J. geol. Soc. Lond.*, **15**, 237-238.
- Guinot, D. (1977). Propositions pour une nouvelle classification des Crustacés Décapodes Brachyours. *C. r. hebd. Séanc. Acad. Sci.*, Paris, (D), **285**, 1049-1052.
- Guinot, D. and M. Taveres (2001), Une nouvelle famille de Crabes du Crétacé, et la notion de Podotremata Guinot, 1977 (Crustacea, Decapoda, Brachyura). *Zoosystema*, **23** (3), 507-546.
- Haan, W. de (1833-50), Crustacea. In Siebold, P. F. von, *Fauna Japonica, sive descriptio animalium, quae in itinere per Japoniam, jussu et auspiciis superiorum, qi summum in India, Batavia Imperium tenet, suscepto anis 1823-30 collegit notis observationibus et adumbrationibus illustravit*. ix-xvi, i-xxxii, vii-xvii + 1-243 p.
- Haj, A. E. and R. M. Feldmann (2002), Functional morphology and taxonomic significance of a novel cuticular structure in Cretaceous raninid crabs (Decapoda: Brachyura: Raninidae). *J. Paleont.*, **76** (3), 472-485.
- Holthuis, L. B. (1949), Fossil decapod Crustacea from the Miocene and younger deposits of the Netherlands. *Med. Geol. Sticht. n. ser.*, **3**, 55-68.
- Holthuis, L. B. (1987), *Necora*, a new genus of European swimming crabs (Crustacea Decapoda, Portunidae) and its type species, *Cancer puber* L., 1767. *Zool. Meded., Leiden*, **61** (1), 1-14.
- Ingle, R. (1980), *British Crabs*. Brit. Mus. (Natural History), London and Oxford University Press, Oxford, 222 p.
- Karasawa, H. and H. Hayakawa (2000), Additions to Cretaceous decapod crustaceans from Hokkaido, Japan – Part 1. Nephropidae, Micheleidae and Galatheididae. *Palaeont. Res.*, **4** (2), 139-145.
- Latreille, P. A. (1802-1803), *Histoire naturelle general et particuliere des crustacés et des insectes*. F. Dufart, Paris, 468 + 391 p.
- Latreille, P. A. (1831), *Cours d'Entomologie ou du l'Histoire naturelle des Crustacés, des arachnides, des Myriapodes et des Insectes: à l'usage des élèves de l'école du Museum d'Histoire naturelle*. Paris, xiii+568+26 p.
- Leach, W. E. (1813-1814), Crustaceology. In Brewster, Sir D. (ed.), *The Edinburgh Encyclopedia: conducted by D. Brewster*. Vol. 7, no. 1, (1813), 383-384; vol. 7, no. 2, (1814), 385-437, Edinburgh.
- Leach, W. E. (1815), A tabular view of the external characters of four classes of animals, which Linné arranged under Insecta, with the distribution of the genera composing three of these classes into orders and descriptions of several new genera and species. *Trans. Linn. Soc., London*, **11**, 306-400.
- Linnaeus, C. (1758), *Systema Naturae per Regna Tria Naturae Secundum Classes, Orcines, Genera, Species, cum Characteribus, Differentiis, Synonymis Locis*. Holmiae. Edition 10, 1, iii+1-578 p.
- Linnaeus, C. (1763), *Systema naturae ...* ed. 12, Tom. 1 *Regnum Animale*. Pt. 2, 533-1327.
- Lockington, W. N. (1876 [1877]), Remarks on the Crustacea of the Pacific Coast of North America, including a Catalogue of Specimens in the Museum of the California Academy of Sciences. *Proc. California Acad. Sci.*, **7**, 63-78.
- Lórenthey, I. (1929), In Lórenthey, I. and K. Beurlen, *Die fossilen Dekapoden der Länder Ungarischen Krone. Geologica hung. Seria palaeontologica*, fasc. 3, 420 p.
- McCoy, F. (1849), On the classification of some British fossil Crustacea with notices of new forms in the University collection at Cambridge. *Ann. Mag. nat. Hist.*, ser. 2, **4**, 161-179, 330-335.
- McCoy, F. (1854), On some new Cretaceous Crustacea. *Ann. Mag. nat. Hist.*, ser. 2, **14**, 116-122.
- MacLeay, W. S. (1838), On the brachyurus Crustacea brought

- from the Cape by Dr Smith. In *Illustrations of the Zoology of South Africa*, 5, *Invertebratae*, 53-71, Smith Elder & Co., London.
- Manning, R. B. (1987), Notes on Western Atlantic Callianassidae (Crustacea: Decapoda Thalassinidae). *Proc. Biol. Soc. Wash.*, **100** (2), 386-401.
- Mantell, G. A. (1844), *Medals of Creation*, 2, i-viii+457-1016, London.
- Mertin, H. (1941), Decapoda Krebse aus dem Subhercynen und Braunschweiger Emscher und Unterseenen sowie Bemerkungen über einige verwante Formen in der Oberkreide. *Nov. Act. Leop.*, NF, **10**, 149-264.
- Meyer, H. von (1854), Jurassische und Triassische Crustaceen. *Palaeontogr.*, **4**, 44 -45.
- Meyer, H. von (1859), *Eryon raiblanus* aus den Raibler-Schichten in Kärnten. *Palaeontogr.*, **8**, 27-30.
- Meyer, H. von (1860), Die Prosoponiden oder Familie der Maskenkrebse. *Palaeontogr.*, **7**, 183-222.
- Milne Edwards, A. (1861), Étude zoologiques sur les Crustacés recents de la famille des Portuniens. *Arch. Mus. Hist. Nat.* (Paris), **10**, 309-428, 2 unnumbered pages.
- Milne Edwards, A. (1880), Étude préliminaires sur les Crustacés, 1ere Partie. In *Reports on the Results of Dredging under the Supervision of Alexander Agassiz, in the Gulf of Mexico and in the Caribbean Sea, 1877, '78, '79, by the U. S. Coast Survey Steamer "Blake", Lieut.-Commander J. R. Bartlett, U. S. N., Commanding*. *Bull. Mus. Comp. Zool. Harvard College*, **8**, 1, 68 p.
- Milne Edwards, H. (1834-37), *Histoire naturelle des Crustacés, comprenant l'anatomie, la physiologie et la classification des ces animaux*. Roret, Paris, 1, xxxv+468 p.; 2, 1-532 p.
- Milne Edwards, H. (1837), *Histoire Naturelle des Crustacés, in L'Institut*, Paris, 225 p.
- Milne Edwards, H. (1850), In d'Archiac, A., *Histoire des Progrés de la Géologie de 1834 à 1849*. 3, Paris.
- Montagu, G. (1808), Description of several marine animals found on the south coast of Devonshire. *Trans. Linn. Soc. Lond.*, **9**, 81-114.
- Montagu, G. (1813), Descriptions of several of new or rare animals, principally marine, discovered on the south coast of Devonshire. *Trans. Linn. Soc. Lond.*, **11**, 1-26.
- Morris, S. F. (1980), Catalogue of the Type and Figured Specimens of Fossil Crustacea (excl. Ostracoda, Chelicerata, Myriapoda and Pycnogonida in the British Museum (Natural History). *Brit. Mus. (Nat. Hist.)*, 52 p.
- Olivi, G. (1782), *Zoologia Adriatica ossia Catalogo ragionato degli animali del Golfo e della Lagune di Venezia; preceduto da una Dissertazione sulla Storia fisica e naturale del Golfo; e accompagnato da Memorie, ed Osservazioni di Fisica Storia naturale ed Economia*. xxxii+334 p.
- Oppel, A. (1862), Ueber Jurassische Crustaceen. *Palaeont. Mitth.* **I**, 120 p.
- Ortmann, A. E. (1892), Die decapoden-Krebse des Strassburger Museum, mit besondere Berücksichtigung der von Herrn. Dr. Döderlein bei Japan und bei den Liu-Kiu-Insel gesammelt und zur Zeit I Strassburger Museum auf bewahrten Formen. *Zool. Jb. (Syst.)*, **6**, 1-58, 241-326, 532-588.
- Ortmann, A. E. (1893), Die Dekapoden-krebse des Strassburger Museums. VII. Theil. Abtheilung; Brachyura genuina Boas) II Unterabtheilung; Cancroidea, 2. Section: Cancrinae. I. Gruppe: Cyclometopa, *Zool. Jb. (Syst.)*, **7**, 411-495.
- Ortmann, A. E. (1898), Arthropoda, In *Dr H. G. Bronn's Klassen und Ordnungen des Thier-Reichs, wissenschaftlich dargestellt in Wort und Bild. Funfter Band. II, Abtheilung. Gliederfussler*. 50-52. C. F. Winter'sche Verlagshandlung.
- Pennant, T. (1777), *Crustacea, Mollusca, Testacea*. *British Zoology*, edition 4, viii+136 p. Penrith, M.-L. and B. F. Kensley.
- Quayle, W. J. (1984), A new crab, *Portunites stintoni* (Crustacea, Decapoda) from the London Clay. *Tert. Res.* **5** (4) 173-178.
- Quayle, W. J. (1987), English Eocene Crustacea (Lobsters and Stomatopod). *Palaeontology*, **30** (3), 581- 612.
- Quayle, W. J. and J. S. H. Collins (1981), New Eocene crabs from the Hampshire Basin. *Palaeontology*, **24**, 733-758.
- Rafinesque Schmaltz, C. (1815), *Analyse de la nature ou tableau de l'univers et de corps organisés*. Palermo, 11-55.
- Rathbun, M. J. (1898), The Brachyura of the biological expedition to the Florida Keys and the Bahamas in 1893. *Bull. Lab. Nat. Hist. State Univ. Iowa*, **4**, 349-375.
- Rathbun, M. J. (1918), The Grapsoid crabs of America. *Bull. U.S. natn. Mus.*, no. 97, xxii+461 p.
- Rathbun, M. J. (1926), The fossil stalk-eyed Crustacea of the Pacific Slope of North America. *Bull. U.S. natn. Mus.*, **138**, 1-155 p.
- Rathbun, M. J. (1930a), The Cancroid crabs of America of the families Euryalidae, Portuni dae, Atelecyclidae, Cancridae, and Xanthidae. *Bull. U.S. natn. Mus.*, **152**, xvi+609 p.
- Rathbun, M. J. (1930b), Fossil decapod crustaceans from Mexico. *Proc. U.S. natn. Mus.*, **78**, 1-10.
- Reuss, A. (1857), Zur Kenntnis fossiler krabben. *Sitzungsber. K. Akad. d. Wissensch., Wien*. **27**, 161-166.
- Saint Laurent, M. de (1980), Sur la classification et la phylogénie des Crustacés Décapodes Brachyours. I. Podotremata Guinot, 1977, et Eubrachyura sect. nov. *C. r. hebd. Séanc. Acad. Sci.*, Paris, (D), **290**, 1265-1268.
- Sakai, K. (1999), Synopsis of the family Callianassidae, with keys to subfamilies, genera and species, and the description of new taxa (Crustacea: Decapoda: Thalassinidea). *Zool. Verhand, Leiden*. **326**, 152 p.
- Salter, J. W. and H. Woodward [Woodward and Salter of some authors] (1865), *Chart of Fossil Crustacea, with*

- descriptive catalogue*. London, Stanford, 1-28.
- Salva, E. W. and R. M. Feldmann (2001), Reevaluation of the family Atelecyclidae (Decapoda: Brachyura). *Kirklandia*, **52**, 9-62.
- Samouelle, G. (1819), *The Entomologist's Useful Compendium, or an Introduction to the knowledge of British Insects*. London, 496 p.
- Sauvage, H. -E. (1891), Note sur les Crustacés des terraines Jurassiques supérieurs du Boulonnais. *An. Sci. Nat. Zool.* 7e sér. **8**, 83-96.
- Schweitzer, C. E. (2000), Tertiary Xanthoidea (Crustacea: Decapoda: Brachyura) from the west coast of North America. *J. Crust. Biol.*, **20** (4), 715-742.
- Schweitzer, C. E. and R. M. Feldmann (2000a), Re-evaluation of the Cancridae Latreille, 1802 (Decapoda: Brachyura) including three new genera and three new species. *Contrib. Zool.*, **69** (4), 223-250.
- Schweitzer, C. E. and R. M. Feldmann (2000b), New species of calappid crabs from western North America and reconsideration of the Calappidae sensu lato. *J. Paleont.*, **74** (2), 230-246.
- Schweitzer, C. E. and R. M. Feldmann (2000c), First notice of the Chirostylidae (Decapoda) in the fossil record and new Tertiary Galatheidae (Decapoda) from the Americas. *Bull. Mizunami Fos. Mus.* **27**, 147-165.
- Schweitzer, C. E. and R. M. Feldmann (2000d), New fossil Portunids from Washington, USA, and Argentina, and a re-evaluation of generic and family relationships within the Portunoidea Rafinesque, 1815 (Decapoda: Brachyura). *J. Paleont.*, **74** (4), 636-653.
- Schweitzer, C. E. and R. M. Feldmann (2001), Differentiation of the fossil Hexpodidae Miers, 1886 (Decapoda: Brachyura) from similar forms. *J. Paleont.*, **75** (2), 330-345.
- Schweitzer, C. E. and E. W. Salva, (2000), First recognition of the Cheiragonidae (Decapoda) in the fossil record and comparison of the family with the Atelecyclidae. *J. Crust. Biol.*, **20** (2), 285-298.
- Simpson, M. (1983), Decapod Crustacea and Associated Fauna. *Proc. Dorset nat. Hist. Archaeol. Soc.*, **104**, 143-146.
- Simpson, M. (1985), The stratigraphy of the Atherfield Clay Formation (Lower Aptian, Lower Cretaceous) of the type and other localities in southern England. *Proc. Geol. Assoc.*, **96** (1), 23-45.
- Stenzel, H. B. (1945), Decapod crustaceans from the Cretaceous of Texas. *Bull. Univ. Texas Bur. Econ. Geol. Technol.*, **4401**, 401-476.
- Stimpson, W. (1858), Prodrum descriptionis anamalium evabratorum, quae in Expeditione ad Oceanum Pacificum Septentrionalem, a Republica Federata mssa, Cadwaladaro Ringgold et Johanne Rogers Ducibus, observavit et descripsit. Crustacea Anomura. *Proc. Acad. Nat. Sci. Philad.*, **10** (7), 225-252.
- Stimpson, W. (1870-1881), Brachyura. Pt. 1. In L. F. de Pourtalés (ed.), *Preliminary report on the Crustacea dredged in the Gulf Stream in the Straits of Florida*. *Bull. Mus. Comp. Zool. Harvard Coll.*, **2**, 150-156.
- Takeda, M. and S. Miyake (1969), A small collection of crabs from New Zealand. *Ohmu. Occ. Pap. Kyushu Univ. Zool. Lab.*, **2**, 157-193.
- Tucker, A. (1998), Systematics of the family Raninidae (Crustacea: Decapoda: Brachyura), with accounts of three new Genera and two new species. *Proc. Biol. Soc. Wash.*, **111** (2), 321-371.
- Van Straelen, V. (1923), Sur les Crustacés décapodes du Bathonien. *Compt. Rend. Acad. Sci. Paris.* **177**, 552-554.
- Van Straelen, V. (1925), Contribution a l'étude des Crustacés décapodes de la période Jurassique. *Mém. Classe des Sci. Acad. r. Belg., Coll. Sci.*, sér. 2, **7**, 461 p.
- Van Straelen, V. (1936), Crustacés décapodes nouveaux ou peu connus de l'époque crétacique. *Bull. Mus. r. Belg. Cl. Sci.*, **12**, 45-50.
- Vega, F. J., R. M. Feldmann, A. C. Ocampo, and K. O. Pope (1997), A new species of late Cretaceous crab (Brachyura: Carcineretidae) from Albion Island, Belize. *J. Paleont.*, **71** (4), 620-637.
- Via Boada, L. (1959), Decápodos fósiles del Eoceno español. *Bol. Inst. Geol. Min. Espana*, **70**, 72 p.
- Wehner, G. (1988), *Über die Prosopiden (Crustacea, Decapoda)*. Inaugural Dissertation, München. 154 p.
- Withers, T. H. (1932), A Liassic crab, and the origin of the Brachyura. *Ann. Mag. nat. Hist.*, ser. 10, **9**, 313-323.
- Withers, T. H. (1951), Some Jurassic and Cretaceous crabs (Prosoponidae). *Bull. nat. Hist. Mus. Lond. (Geol.)*, **1** (6), 172-192.
- Woods, H. (1925-31), *A Monograph of the Fossil Macrurous Crustacea of England*. Palaeontogr. Soc. [Monogr.], 1-122 p.
- Woods, J. T. (1953), Brachyura from the Cretaceous of central Queensland. *Mem. Qd. Mus.*, **13**, 50-57.
- Woodward, H. (1866), On the oldest known British crab (*Palaeinachus longipes*) from the Forest Marble of Malmesbury. *Q. J. Geol. Soc. Lond.*, **5**, 258-261.
- Woodward, H. (1867), On a new genus of shore crab, *Goniocypoda Edwardsi*, from the Lower Eocene of Hampshire. *Geol. Mag.*, **4**, 529-531.
- Woodward, H. (1868), On a new brachyurous crustacean (*Prosopon mammillatum*) from the Great Oolite, Stonesfield. *Geol. Mag. Lond.*, **5**, 3-5.
- Woodward, H. (1869), Fourth Report on the Structure and Classification of the fossil Crustacea. *Report of the 38th Meet. Brit. Assoc. Adv. Sci.* (Norwich 1868), 72-75.
- Woodward, H. (1871), On some new Lower Eocene crustaceans from Portsmouth. *Q. J. Geol. Soc. Lond.*, **27**, 90-92.
- Woodward, H. (1873), Further notes on Eocene Crustacea from Portsmouth. *Q. J. Geol. Soc. Lond.*, **29**, 25-31.
- Woodward, H. (1876), On some new macrurous Crustacea from the Kimmeridge Clay of the Sub-Wealden Boring, Sussex, and from Boulogne-sur-Mer. *Q. J. Geol. Soc. Lond.*, **32**, 47-50.
- Woodward, H. (1907), On a new brachyurus Crustacean from

- (the "Clypeus Grit" (Inf. Oolite) of the Cotteswold Hills. *Geol. Mag.*, N. S. dec. 5, **8**, 307-311.
- Wright, C. W. (1997), New information on Cretaceous crabs. *Bull. nat. Hist. Mus. Lond. (Geol.)*, **53** (2), 135-138.
- Wright, C. W. and J. S. H. Collins (1972), *British Cretaceous crabs*. Palaeont. Soc. Monogr., no. 126, 114 p.
- Wright, C. W. and J. S. H. Collins (1975), *Glaesnerella* (Crustacea Decapoda Cymonomonidae), a replacement name for *Glaessneria* Wrght and Collins, 1972 *non Glaessneria* Takeda and Miyake, 1969. *Palaeontology*, **18** (2), 441.
- Zariquiey Alvarez, R. (1946), *Crustáceos decápodos Mediterráneos*. *Inst. Españ. Estud. Medit. Publ. Biol. Medit.*, 181 p.

Manuscript accepted on August 29, 2002